

**MOUNTAIN SERIES WEEKENDER MEDICAL - benzalkonium chloride, povidone-iodine, acetaminophen, aspirin, diphenhydramine hydrochloride, ibuprofen, bacitracin zinc, neomycin sulfate, polymyxin b sulfate**

**Tender Corporation dba Adventure Medical Kits**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Mountain Series Weekender Medical Kit**

**Active Ingredient**

Active Ingredient:

Benzalkonium Chloride 0.40%

**Purpose**

Antiseptic

**Use**

For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.

**Warnings**

Warning: For external use only.

Keep out of reach of children: If swallowed, get medical help or contact a Poison Control Center right away.

Stop use if unusual redness, swelling or other symptoms occur. Consult a physician immediately.

Do not use in the eyes or over large areas of the body.

**Directions**

Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.

**Inactive Ingredients**

Inactive Ingredient: Purified water

LOT/EXP:                      Made in CHINA

20130301

Antiseptic Towelette

Genuine First Aid LLC, Clearwater FL 33755

www.GenuineFirstAid.com

1/pouch

GENUINE FIRST AID

### **Active Ingredients**

Active Ingredient: .....Bacitracin Zinc 400 units

Neomycin Sulfate 5mg ( equivalent to 3.5 mg Neomycin base)

Polymyxin B Sulfate 5000 units

### **Purpose**

Triple Antibiotic

Uses: To help prevent infection in:  
minor cuts; scrapes; burns

### **Warnings**

For external use only.

Do not use: in eyes; over large areas of the body;

If allergic to any of the ingredients; for more than one week unless directed by a physician.

Stop use and consult a doctor:

if the condition persists or gets worse; a rash or other allergic reaction develops

Keep out of reach of children.

If ingested, contact a Poison

Control Center right away.

### **Directions**

Directions: clean affected area; apply small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily; may be covered with a sterile bandage

Other information:

Store at room temperature.

### **Inactive Ingredient**

water

Genuine Triple Antibiotic

First Aid Ointment

To Help Prevent Infection

Each Gram Contains:

Bacitracin Zinc 400 units

Neomycin Sulfate 5 mg

(equivalent to 3.5 mg

Neomycin base)

Polymyxin B Sulfate 5000 units

Net Wt. 0.5g ; (1/64 oz)

Manufactured in CHINA for

GENUINE FIRST AID.

Triple Antibiotic Ointment 10pcs

Net wt. 0.9g (1/32oz)

100

Triple Antibiotic

### **Active ingredient (in each tablet)**

Aspirin (NSAID\*) 325 mg

\*nonsteroidal anti-inflammatory drug

### **Purpose**

Pain reliever/fever reducer

### **Uses**

**Temporarily relieves minor aches and pains associated with**

- headache
- minor arthritis
- common cold
- menstrual cramps
- muscular aches
- backache
- toothache

**Temporarily reduces fever.**

### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcohol drinks every day while using this product
- take more or for a longer time than directed

#### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/ fever reducer
- right before or after heart surgery
- if you are taking prescription drugs for gout, diabetes or arthritis

#### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

#### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

#### **When using this product**

- take with food or milk if stomach upset occurs

#### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - you have difficulty swallowing
  - if ringing in the ears or loss of hearing occurs redness or swelling is present in the painful area
  - any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- drink a full glass of water with each dose

**Adults and children: (12 years and older)**

Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

**Children under 12 years:**

Do not give to children under 12 years of age.

**Other information**

- read all product information before using
- store at room temperature 59°-86°F (15°-30°C)
- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

**Inactive ingredients**

croscarmellose sodium\*, hypromellose, microcrystalline cellulose\*, mineral oil\*, polyethylene glycol\*, propylene glycol\*, starch, titanium dioxide\*

*\*may contain*

**Questions or comments?**

1-800-634-7680

**116R Medique APAP 325 mg Label**

Medique®

Aspirin

5 Grain (325 mg)

Pain Reliever/Fever Reducer (NSAID)

See new warnings information

Easy To Swallow

Film Coated Tablets

24 Tablets

(12 x 2)

Compare active ingredients to Bayer®

Registered Trademark Bayer Consumer

Tamper Evident Unit Dose Packets

**Active ingredients**

(in each capsule)

Diphenhydramine Hydrochloride 25 mg

**Purpose**

Antihistamine

## Uses

Temporarily relieves runny nose and decreases sneezing, itching of the nose and throat, itchy-watery eyes due to hay fever or other respiratory allergies.

## Warnings

### Do not use

- with any other product containing diphenhydramine, even one that is used on skin.

### Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- glaucoma

### Ask a doctor or pharmacist before use if you are

- taking any drugs for asthma
- taking sedatives or tranquilizers

### When Using This Product

- marked drowsiness may occur
- avoid alcohol beverages
- alcohol, sedatives and tranquilizers may increase the drowsiness effect
- use caution when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**If pregnant or breast feeding,** ask a health professional before use.

**Keep out of the reach of children.** In case of overdose, contact a physician or poison control center immediately.

## Directions

- **do not use more than directed**

**Adults and children: (12 years and older):** Take 1 to 2 capsules every 4 to 6 hours as needed. Do not take more than 12 capsules in 24 hours.

**Children under 12 years:** Do not give to children under 12 years of age.

## Other information

- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

## Inactive ingredients

D&C Red #28, FD&C Blue #1, FD&C Red #40, gelatin, starch

**Questions or comments? 1-800-634-7680**

## **182R Medique Diphen Label**

Collect Medi-Bucks

See inside flap for more details

Medique®

Diphen

Diphenhydramine HCl 25 mg

Hay Fever/Allergies

Fiebre del Heno/Alergias

Pull to Open

TiraParaAbrir

Easy To Swallow Capsules

Capsulas Faciles de Tragar

200 Capsules

(200 x 1)

Tamper Evident Unit Dose Packets

Empaquetado con Sellado Evidente en Dosis Unitarias

### ***Active ingredient (in each tablet)***

Ibuprofen (NSAID\*) 200 mg

\*nonsteroidal anti-inflammatory drug

### ***Purpose***

Pain reliever/fever reducer

### ***Uses***

**Temporarily relieves minor aches and pains associated with**

- headache
- backache
- common cold
- minor arthritis pain
- toothache
- menstrual cramps
- muscular aches

**Temporarily reduces fever.**

### ***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction which may include:

- hives
- asthma (wheezing)
- rash
- blisters
- skin reddening

- facial swelling
- shock

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts for more than 10 days
  - fever gets worse or lasts for more than 3 days
  - redness or swelling is present in the painful area
- any new or unexpected symptoms occur

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless specifically directed to so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- **do not use more than directed**
- **the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see Warnings)

**Adults and children: (12 years and older)**

Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

**Children under 12 years:**

Do not give to children under 12 years of age.

**Other information**

- read all product information before using
- store at 68-77°F (20-25°C)
- avoid excessive heat 40°C (above 104°F)
- tamper evident sealed packets
- do not use any opened or torn packets

**Inactive ingredients**

carnauba wax\*, cellulose\*, colloidal silicon dioxide, corn starch\*, hypromellose, iron oxide red\*, lactose, magnesium stearate, microcrystalline cellulose\*, polydextrose, polyethylene glycol, povidone, silica\*, sodium lauryl sulfate\*, sodium starch glycolate, stearic acid\*, titanium dioxide, triacetin\*

\* may contain

**Questions or comments?** 1-800-634-7680

**Principal Display Panel**

Medi-First®

Ibuprofen 200 mg

Pain Reliever/Fever Reducer (NSAID)

Easy To Swallow Film Coated Tablets

See new warnings information

100 tablets (50 x 2)

Compare active ingredient to: Genuine Advil®

Registered trademark Wyeth Consumer

Tamper Evident Unit Dose Packets

**Active ingredient (in each tablet)**

Acetaminophen 325 mg

**Purpose**

Pain reliever/fever reducer

**Uses****For the temporary relief of minor aches and pains associated with**

- headache
- common cold
- muscular aches
- toothache
- minor arthritis pain
- menstrual cramps

## **For the reduction of fever.**

### **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

### **Ask a doctor before use if you have**

- liver disease

### **Ask a doctor or pharmacist before use if**

- you are taking the blood thinning drug warfarin

### **Stop using and ask a doctor if**

- symptoms do not improve
- new symptoms occur
- pain or fever persists or gets worse
- redness or swelling is present

**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**If pregnant or breast-feeding,** ask a health professional before use.

### **Directions**

- **do not use more than directed**

### **Adults and children: (12 years and older)**

Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.

### **Children under 12 years:**

Do not give to children under 12 years of age.

### **Other information**

- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

### **Inactive ingredients**

corn starch, hypromellose, maltodextrin\*, microcrystalline cellulose\*, polyethylene glycol, povidone, pregelatinized starch\*, sodium starch glycolate, stearic acid

\* *may contain*

**Questions or comments? 1-800-634-7680**

## **Principal Display Panel**

### **145R Medique APAP 325 mg Label**

Collect MediBucks

See inside flap for more details

Medique®

APAP

Acetaminophen 325 mg

Pain Reliever/Fever Reducer

Alivia el Dolor/Reduce la Fiebre

Easy To Swallow

Film Coated Tablets

Facil de Tragar Tabletillas con Cubierta Pelicular

Pull to Open

Tire Para Abrir

See new warnings information

500 Tablets

(250 x 2)

Tamper Evident Unit Dose Packets

Empaquetado con Sellado Evidente en Dosis Unitarias

### **Weekender Kit Label**

Mountain Series Medical Kit Weekender

Adventure Medical Kits BE SAFE

1-6 People up to 7 days

Povidone-iodine 10%

Antiseptic

Warnings

Do not use

- if allergic to iodine
- in the eyes

For external use only

Ask a doctor before use if injuries are

- deep or puncture wounds
- serious burns

Stop use and ask a doctor if

- redness, irritation, swelling or pain persists or increases
- infection occurs

Avoid pooling beneath patient

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or consult a poison control center immediately.

NDC 52380-0001-1  
NPN 02076144



**ONE FLUID OUNCE**  
**POVIDONE-IODINE**  
**SOLUTION**

1 Fl. oz. (30 mL)

**STERILE** unless opened or damaged.

<b>Drug Facts</b>	
<b>Active ingredient</b>	<b>Purpose</b>
Povidone-iodine USP 10%.....	Antiseptic
<b>Use</b> antiseptic skin preparation	
<b>Warnings</b>	
Do not use ■ if allergic to iodine ■ in the eyes	
For external use only ▶	

Reorder No. L-1001



LOT



**Drug Facts** (continued)

**Warnings**

Ask a doctor before use if injuries are  
■ deep or puncture wounds ■ serious burns

**Stop use and ask a doctor if**

- redness, irritation, swelling or pain persists or increases
- infection occurs

Avoid pooling beneath patient

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or consult a poison control center immediately.

**Directions** apply locally as needed

**Other information**

- 1% titratable iodine ■ latex free
- for hospital or professional use only

**Inactive ingredients** citric acid, disodium phosphate, nonoxonyl-9, sodium hydroxide, water

**Questions or comments?**

☎ 1 800 760-3236 (Mon to Fri 8:30 AM-5:00 PM EST)



(01) 0 0352380 0001 9

**APLICARE, INC.**  
**MERIDEN, CT 06450**  
**U.S.A.**  
DOLLARD-des-ORMEAUX,  
QUEBEC H9G 2S3  
www.aplicare.com  
0609

Medique®

2 Tablets

Made in USA

APAP

Active ingredient (in each tablet) Purpose  
Acetaminophen 325 mg..... Pain reliever/  
fever reducer

Uses

For the temporary relief of minor aches and pains associated with

- headache
- muscular aches
- minor arthritis pain
- toothache
- common cold
- menstrual cramps

For the reduction of fever.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Store at 59°-86°F (15-30°C) • Tamper-evident sealed packets • Do not use any opened or torn packets  
Mfd. for MEDIQUE PRODUCTS • Fort Myers, FL 33967

Warnings (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if you have

- liver disease

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin

Stop using and ask a doctor if

- symptoms do not improve
- new symptoms occur
- pain or fever persists or gets worse
- redness or swelling is present

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

If pregnant or breast-feeding, ask a health professional before use.

Directions ■ do not use more than directed

Adults and children: (12 years and older)

Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.

Children under 12 years:

Do not give to children under 12 years of age.

SEE CARTON FOR COMPLETE PRODUCT INFORMATION

Lot 1831  
Exp. 01-2013

Made in  
**USA**

2 Tablets

# IBUPROFEN

**Active ingredient (in each tablet)**      **Purpose**

Ibuprofen (NSAID\*) 200mg ..... Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** Temporarily relieves minor aches and pains due to

- headache ■ toothache ■ backache ■ menstrual cramps
- common cold ■ muscular aches ■ minor arthritis pain

Temporarily reduces fever.

## Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction which may include: ■ hives ■ skin reddening ■ facial swelling ■ rash ■ asthma (wheezing) ■ blisters ■ shock If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older ■ have stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Store at 68-77°F (20-25°C) • Avoid excessive heat, above 104°F (40°C) • Tamper-evident sealed packets

• Do not use any opened or torn packets

Mfd. for **MEDIQUE PRODUCTS** • Fort Myers, FL 33967

## Warnings (continued)

### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

### Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

### When using this product

- take with food or milk if stomach upset occurs

### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint ■ vomit blood ■ have bloody or black stools
  - have stomach pain that does not get better

- pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## Directions

- do not use more than directed
- the smallest effective dose should be used

**Adults and Children (12 years and older):** Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

**Children under 12 years:** Do not give to children under 12 years of age.

Lot 1922 Exp. 02-2012

2 Tablets

# Medique<sup>®</sup> Aspirin

Made In  
USA

### Active ingredient (in each tablet)

Aspirin (NSAID\*) 325 mg  
\*nonsteroidal anti-inflammatory drug

### Purpose

Pain reliever/fever reducer

**Uses** Temporarily relieves minor aches and pains due to  
■ headache ■ muscular aches ■ minor arthritis pain ■ backache  
■ common cold ■ toothache ■ menstrual cramps  
Temporarily reduces fever.

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ skin reddening ■ facial swelling ■ rash ■ asthma (wheezing) ■ blisters ■ shock  
If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
■ are age 60 or older ■ have had stomach ulcers or bleeding problems  
■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Store at 59°-86°F (15°-30°C) • Tamper-Evident Sealed Packets  
• Do Not Use any Opened or Torn Packets

Mfd. for **MEDIQUE PRODUCTS** • Fort Myers, FL 33967

### Warnings (continued)

Do not use

■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery ■ if you are taking a prescription drug for gout, diabetes or arthritis

Ask a doctor before use if

■ stomach bleeding warning applies to you  
■ you have a history of stomach problems such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are

■ under a doctor's care for any serious condition ■ taking any other drug

When using this product

■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:  
■ feel faint ■ vomit blood ■ have bloody or black stools  
■ have stomach pain that does not get better  
■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ you have difficulty swallowing ■ if ringing in the ears or loss of hearing occurs ■ redness or swelling is present in the painful area  
■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions:** Do not use more than directed

**Adults and children: (12 years and older)** Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

**Children: (under 12 years)** Do not give to children under 12 years of age.

SEE CARTON FOR COMPLETE PRODUCT INFORMATION

Lot 1856 Exp. 11-2012

**Medique**<sup>®</sup>

NDC# 47682-182-46

Made in  
USA

**Diphen**  
1 Capsule

**Active Ingredient**

(In each capsule)

Diphenhydramine

Hydrochloride 25mg .....Antihistamine

**Purpose**

**Uses:**

Temporarily relieves the following symptoms associated with hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes

Store at room temperature • Tamper-Evident Sealed  
Packets • Do Not Use any Opened or Torn Packets

Mfd. for **MEDIQUE PRODUCTS**

Fort Myers, FL 33967

**Warnings**

Ask a Doctor Before Use

If you have:

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

If you are:

- taking any drugs for asthma
- taking sedatives or tranquilizers

When Using this product:

- drowsiness may occur
  - avoid alcohol beverages
- alcohol, sedatives and tranquilizers may increase the drowsiness effect

use caution when driving a motor vehicle or operating machinery  
excitability may occur, especially in children.

do not exceed recommended dosage. Keep this and all drugs  
out of the reach of children. In case of accidental overdose,  
contact a physician or poison control center immediately. As with  
any drug, if you are pregnant or nursing a baby, seek the advice  
of a health professional before using this product.

**Directions:** Do not use more than directed

Adults and children: (12 years and older) Take 1 capsule  
every 4 to 6 hours as needed. Do not take more than 12 capsules  
in 24 hours, or as directed by a doctor.

Children under 12 years: Do not give to children under 12 years  
of age unless directed by a doctor.

Lot 1555 Exp. 1/2010

REORDER AST-001

**Antiseptic  
Towelette**  
**Toallitas  
Antisepicas**



Genuine First Aid LLC, Clearwater FL 33755  
www.GenuineFirstAid.com



**GENUINE FIRST AID®**



DRUG FACTS - Antiseptic Towelette

Active Ingredient: Benzalkonium Chloride 0.40% Purpose: First Aid Antiseptic

Use: For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.

Warnings: For external use only. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. If unusual redness, swelling or other symptoms occur, consult a physician immediately.

Do not use: In the eyes, or over large areas of the body.

Directions: Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.

Inactive ingredient: Purified water

Made in CHINA

LOT/EXP:  
20130301

--- TEAR HERE ---

Reorder TAO-001

# Genuine Triple Antibiotic

**First Aid Ointment  
To Help Prevent Infection**

**Each Gram Contains:  
Bacitracin Zinc 400 units  
Neomycin Sulfate 5 mg  
(equivalent to 3.5 mg  
Neomycin base)  
Polymyxin B Sulfate 5000 units  
Net Wt 0.5g • (1/64 oz)**



Manufactured in CHINA for  
**GENUINE FIRST AID.**

## Drug Facts

Uses: To help prevent infection  
in • minor cuts • scrapes • burns

## Warnings:

For external use only

Do not use • in eyes • over large  
areas of the body • if allergic to  
any of the ingredients • for more  
than one week unless directed  
by a physician

Stop use and consult a doctor  
• if the condition persists or gets  
worse • a rash or other allergic  
reaction develops

Keep out of reach of children.

If ingested, contact a Poison  
Control Center right away.

Directions: • clean affected area  
• apply small amount of this product  
(an amount equal to the surface  
area of the tip of a finger) on the  
area 1 to 3 times daily • may be  
covered with a sterile bandage

Other information:

Store at room temperature

LOT/EXP:

20130118



ACTIVITY

1-6 PEOPLE  
UP TO 7 DAYS


Adventure  
Medical  
Kits

BE SAFE

MOUNTAIN SERIES

# MEDICAL KIT

WEEKENDER



## MOUNTAIN SERIES WEEKENDER MEDICAL

benzalkonium chloride, povidone-iodine, acetaminophen, aspirin, diphenhydramine hydrochloride, ibuprofen, bacitracin zinc, neomycin sulfate, polymyxin b sulfate kit

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:44224-0118
---------------------	----------------	---------------------------	----------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:44224-0118-1	1 in 1 KIT		

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	6 PACKAGE	4.8 mL
Part 2	3 TUBE	1.5 g
Part 3	2 PACKET	4

<b>Part 4</b>	2 PACKET	2
<b>Part 5</b>	1 PACKET	2
<b>Part 6</b>	2 PACKET	4
<b>Part 7</b>	1 PACKET	22 g

## Part 1 of 7

### ANTISEPTIC TOWELETTE

benzalkonium chloride swab

#### Product Information

**Item Code (Source)** NDC:52124-0001

**Route of Administration** TOPICAL

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.40 mL in 100 mL

#### Inactive Ingredients

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0001-1	0.8 mL in 1 PACKAGE		

#### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	04/23/2010	

## Part 2 of 7

### GENUINE TRIPLE ANTIBIOTIC

bacitracin zinc, neomycin sulfate, polymyxin b sulfate ointment

#### Product Information

**Item Code (Source)** NDC:52124-0003

**Route of Administration**

TOPICAL

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN ZINC	400 [iU] in 1 g
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:16QD7X297)	NEOMYCIN SULFATE	5 mg in 1 g
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B SULFATE	5000 [iU] in 1 g

**Inactive Ingredients**

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0003-1	0.5 g in 1 TUBE		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333B	02/16/2010	

**Part 3 of 7****MEDIQUE ASPIRIN**

aspirin tablet, film coated

**Product Information**

Item Code (Source)	NDC:47682-116
Route of Administration	ORAL

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

**Inactive Ingredients**

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

STARCH, CORN (UNII: O8232NY3SJ)

### Product Characteristics

Color	white (white)	Score	no score
Shape	ROUND (ROUND)	Size	10mm
Flavor		Imprint Code	44;157;aspirin
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-116-99	2 in 1 PACKET		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	12/30/2008	

## Part 4 of 7

### MEDIQUE DIPHEN

diphenhydramine hydrochloride capsule

### Product Information

Item Code (Source)	NDC:47682-182
Route of Administration	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg

### Inactive Ingredients

Ingredient Name	Strength
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN (UNII: 2G866QN327L)	
STARCH, CORN (UNII: O8232NY3SJ)	

### Product Characteristics

<b>Color</b>	pink (pink) , white (white)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (CAPSULE)	<b>Size</b>	14mm
<b>Flavor</b>		<b>Imprint Code</b>	CPC;835
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-182-46	1 in 1 PACKET		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	12/30/2008	

### Part 5 of 7

### MEDI-FIRST IBUPROFEN

ibuprofen tablet, film coated

### Product Information

<b>Item Code (Source)</b>	NDC:47682-808
<b>Route of Administration</b>	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

### Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

## Product Characteristics

Color	white (white)	Score	no score
Shape	ROUND (ROUND)	Size	10mm
Flavor		Imprint Code	44;352
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-808-99	2 in 1 PACKET		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075139	12/30/2008	

## Part 6 of 7

### MEDIQUE APAP

acetaminophen tablet, film coated

## Product Information

Item Code (Source)	NDC:47682-145
Route of Administration	ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

## Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

## Product Characteristics

Color	white (white)	Score	no score
-------	---------------	-------	----------

<b>Shape</b>	ROUND (ROUND)	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	AZ;234
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-145-99	2 in 1 PACKET		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	12/30/2008	

### Part 7 of 7

### APLICARE POVIDONE-IODINE SOLUTION

povidone-iodine solution solution

### Product Information

<b>Item Code (Source)</b>	NDC:52380-0001
<b>Route of Administration</b>	TOPICAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POVIDONE-IODINE (UNII: 85H0HZU99M) (IODINE - UNII:9679TC07X4)	POVIDONE-IODINE	9.8 g in 100 g

### Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM PHOSPHATE, DIBASIC (UNII: GR686LBA74)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
NONOXYNOL-9 (UNII: 48Q180SH9T)	
WATER (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52380-0001-3	22 g in 1 PACKET		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	03/01/1984	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333A	08/01/2011	

**Labeler** - Tender Corporation dba Adventure Medical Kits (064437304)

Revised: 8/2011

Tender Corporation dba Adventure Medical Kits